

PRODUCT INFORMATION SHEET

Information Provided for Australian Veterinary Professionals Only

ilium
SYNTOCIN
INJECTION OF
SYNTHETIC OXYTOCIN



WHAT IS ILIUM SYNTOCIN INJECTION?

- Indications:** For stimulation of uterine contractions and as an aid in the control of minor postpartum haemorrhage and in the removal of retained placenta in horses, cattle, sheep, pigs, dogs and cats.
- Active Ingredients:** 10 IU/mL oxytocin
- Schedule:** S4
- Physical Description:** Clear colourless liquid

WHY CHOOSE ILIUM SYNTOCIN INJECTION?

- ✓ 7-week in-use period following first broaching may help reduce product wastage.
- ✓ Available in 50 mL sterile multi-dose vial.
- ✓ Australian made.
- ✓ Australian owned.
- ✓ Available from your preferred wholesaler.

HOW TO USE ILIUM SYNTOCIN INJECTION?

Administration Route: Intravenous, intramuscular or subcutaneous injection.

Dose Rate:

Horses and cattle: 1–4 mL, S.C., I.M. or I.V.
Sheep and pigs: 0.2–1 mL, S.C., I.M. or I.V.
Dogs and cats: 0.2–1 mL, S.C., I.M. or I.V.

Withholding Periods: MEAT: Zero (0) days. MILK: Zero (0) days.

EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product. Note—observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Troy Laboratories on (+61) 2 8808 3600 before using this product.

Storage Conditions: Store between 2 °C and 8 °C (refrigerate, do not freeze). Protect from light. Use contents within 7 weeks (49 days) of first broaching the vial. Discard the unused portion.

Please refer to carton for full product information.

