


PRODUCT INFORMATION SHEET

Information Provided for Australian Veterinary Professionals Only

ilium
ATIPAMEZOLE
INJECTION



WHAT IS ILIUM ATIPAMEZOLE INJECTION?

Indications:	Reversal and abolishment of effects of medetomidine hydrochloride in cats and dogs.
Active Ingredient:	5 mg/mL atipamezole hydrochloride
Schedule:	S4
Physical Description:	Clear, colourless solution

WHY CHOOSE ILIUM ATIPAMEZOLE INJECTION?

- ✓ 90-day or 3 month in-use broach period.
- ✓ Practical 10 mL multi-dose vial.
- ✓ Predictable reversal when medetomidine sedation is no longer required.
- ✓ Easy dose calculation based on the medetomidine dose already given.
- ✓ Suitable for use in both dogs and cats.
- ✓ Australian made and Australian owned.
- ✓ Conveniently available through your preferred wholesaler.

HOW TO USE ILIUM ATIPAMEZOLE INJECTION?

Dose Rate: Administer by intramuscular injection. ilium Atipamezole Injection is quickly absorbed and is generally administered 15 to 60 minutes after the medetomidine dose.

Dogs: Give a dose five times that of the previous medetomidine dose (50 to 400 µg/kg); by quantity, the same volume as a medetomidine 1 mg/mL injection dose.

Cats: Give a dose 2.5 times that of the previous medetomidine dose (125 to 375 µg/kg); by quantity, half the volume of a medetomidine 1 mg/mL injection dose.

Administration Route: Intramuscular injection.

Contraindications: Do not administer to pregnant bitches.

Precautions: Due to the potent pharmacological activity of atipamezole, skin contact with ilium Atipamezole Injection should be avoided. Wipe any excessive spillage from the animal's skin.

Side Effects: In dogs, a transient hypotensive effect has been observed during the first 10 minutes after injection. Vomiting and panting after atipamezole injection have been reported, but these effects appear to be rare. Transient over-alertness and tachycardia may be observed after a possible overdose.

Storage Conditions: Store below 30 °C (room temperature). Protect from light. Store the vial upright. Use the contents of this vial within 90-days or 3 months of initial broaching. Discard any unused portion.

Please refer to the approved product label for full product information.