

AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

ilium Oxytet-200 L.A.

LONG-ACTING BROAD-SPECTRUM
ANTIBIOTIC INJECTION



WHAT IS ILIUM OXYTET-200 L.A. LONG-ACTING BROAD-SPECTRUM ANTIBIOTIC INJECTION?

ilium Oxytet-200 L.A. long-acting broad-spectrum antibiotic injection is for the treatment and control of conditions caused by oxytetracycline-sensitive organisms in cattle, sheep and pigs.

Active Ingredient: 200 mg/mL oxytetracycline

Poison Schedule: S4

Physical Description: Pale yellow to brown liquid.

WHY CHOOSE ILIUM OXYTET-200 L.A. LONG-ACTING BROAD-SPECTRUM ANTIBIOTIC INJECTION?

- ✓ **Long-acting** and provides a broad-spectrum of treatment.
- ✓ **Targets a large number of Gram-positive and Gram-negative bacteria** – *Mycoplasma* spp., *Chlamydia* and *Rickettsiae*.
- ✓ **Treats multiple clinical conditions** including gastrointestinal tract infections, respiratory infections, urogenital infections, septicaemia and infected wounds.
- ✓ Specific conditions may include:
 - **Cattle:** Anaplasmosis, blackleg, diphtheria, actinobacillosis, pneumonia, footrot, systemic mastitis, joint-ill, navel-ill and leptospirosis.
 - **Pigs:** Systemic mastitis, erysipelas, joint-ill, navel-ill, pneumonia and leptospirosis. An aid in the control of MMA (Metritis-Mastitis-Agalactia) syndrome in sows.
 - **Sheep:** Footrot, systemic mastitis, metritis, blackleg, joint-ill, navel-ill and pneumonia.
- ✓ **Available in 100 and 250 mL sizes.**
- ✓ **6-month broach period**, which allows for plenty of time to use the product and reduces the chance that product will be wasted.
- ✓ **Plastic packaging** which reduces the chance of breakage.
- ✓ **Australian made.**

AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only


ilium Oxytet-200 L.A.

LONG-ACTING BROAD-SPECTRUM
ANTIBIOTIC INJECTION



HOW TO USE ILIUM OXYTET-200 L.A. LONG-ACTING BROAD-SPECTRUM ANTIBIOTIC INJECTION?

Dose Rate:

	DOSE RATE	ADMINISTRATION ROUTE
 CATTLE, SHEEP AND PIGS	1 mL / 10 kg body weight	Deep I.M. only. Production animals should be injected high in the neck to avoid carcass damage.

- No more than 10 mL should be injected into any one site in cattle and no more than 5 mL injected into any site in pigs and sheep.
- A single treatment is usually sufficient. In severe cases a second dose may be administered 72 hours after the initial dose if necessary to achieve the desired clinical response.
- Treated animals should be clearly identified in such a way that they will maintain their identity during the withholding period.
- This preparation may darken during use but potency will remain unaffected.

Side Effects:

- Product may cause anaphylaxis in cattle. Adrenaline may be a suitable antidote. Some tissue reaction and discomfort may occur at injection site.

Contraindications:

Contraindicated for use in horses, dogs and cats.

Withholding Periods:

- MEAT: 42 days.
- MILK: 7 days. Do not feed treated milk to bobby calves.

Trade Advice - Export Slaughter Interval (ESI):

An ESI has not been established for this product. Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may result in the need to extend the approved withholding period.

Storage Conditions:

Store below 30 °C (room temperature). Protect from light.

Please read carton and/or leaflet for full product information.