

ILIMUM BUCCALGESIC® OTM



WHAT IS ILIMUM BUCCALGESIC OTM?

Ilium Buccalgesic OTM Pain Relief for Cattle and Sheep is a non-steroidal anti-inflammatory drug (NSAID) for the alleviation of pain associated with routine husbandry procedures in cattle and sheep.




Active Ingredient: 10 mg/mL meloxicam

Schedule: S6

Physical Description: Blue viscous gel

WHY CHOOSE ILIMUM BUCCALGESIC OTM?

- ✓ **Simply apply blue gel** between the molar teeth and inside of the cheek.
- ✓ **Safe delivery with no needles.**
- ✓ **Long-lasting pain relief** that starts working within 15 minutes.^{1,2,3}
- ✓ **Use for up to 6 months** after opening.
- ✓ **Cost-effective** aid in alleviating pain associated with routine husbandry procedures.
 - **Cattle:** Disbudding (or dehorning) and castration in calves. It is recommended Ilium Buccalgesic is used in combination with a cornual nerve block and local anaesthetic at the surgical site to enhance pain relief and minimise tissue damage and distress.
 - **Sheep:** Castration, tail docking and mulesing of lambs.
- ✓ **Available in 450 mL.**
- ✓ **Developed and made in Australia** in association with






References:

1. Colvin, A. October 2002. Trends in mulesing, tail docking and castration practices of Australian woolgrowers: Results of the 2021 AWI Merino Husbandry Practices Survey. AWI Project No.: ON-00829.
2. Van der Saag, D; Lomax, S; Windsor, P. A.; Taylor, C; Thomson, P; Hall, E; and Whit, P.J. 2018. Effects of topical anaesthetic and buccal meloxicam on average daily gain, behaviour and inflammation of unweaned beef calves following surgical castration. Animal 2018 Nov;12(11):2373-2381.
3. Troy Animal Healthcare Data on File.

ILIMUM BUCCALGESIC®

OTM



HOW TO USE ILIMUM BUCCALGESIC OTM?

Dose Rate:

ANIMAL	DOSE	ILIMUM BUCCALGESIC APPLICATOR GUN AND NOZZLE
<p>CATTLE</p>	<p>0.5 mg/kg body weight (0.5 mL solution /10 kg)</p>	
<p>SHEEP</p>	<p>1.0 mg/kg body weight (1.0 mL solution /10 kg)</p>	

Administration Route:

Administer the dose immediately before the procedure into the space between the molar teeth and the inside of the cheek while holding the mouth closed. This product is not intended to be swallowed (although will not cause harm if this occurs) but to be absorbed through the mucosa of the inner cheek.

Storage Conditions:

Store below 30 °C (room temperature). Protect from light. Viscosity of the product can increase in cold weather. Gently warm bottle in warm water prior to use.

Overdose: In case of overdose, seek veterinary advice. Symptomatic treatment should be initiated.

Restraints:

- DO NOT USE in lactating or pregnant cows where milk may be used or processed for human consumption.

Retreatment Interval:

- **Cattle:** DO NOT re-treat animals for 21 days after treatment.
- **Sheep:** This product is indicated for a SINGLE administration at the recommended dose rate.

Contraindications:

- Contraindicated for use in animals suffering from haemorrhagic gastrointestinal disorders, impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of individual hypersensitivity to the product.
- This product is contraindicated for concurrent administration with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents.

Withholding Periods:

- MEAT: Cattle: 14 days. Sheep: 10 days.
- MILK: Do not use in lactating ewes or pregnant ewes within 10 days of lambing where milk may be used or processed for human consumption.

TRADE ADVICE - Export Slaughter Interval (ESI): Cattle: 21 days. Sheep: 10 days.

Please read carton for full product information.