

AU PRODUCT INFORMATION SHEET



WHAT IS TROY DORAMATE INJECTABLE ENDECTOCIDE?

Troy Doramate Injectable Endectocide is a broad-spectrum parasiticide for the treatment and control of doramectin-sensitive internal and external parasites of cattle and pigs.

Physical Description: Clear, colourless to pale yellow solution.

Active Ingredient: 10 mg/mL DORAMECTIN

WHY CHOOSE TROY DORAMATE INJECTABLE ENDECTOCIDE?

- ✓ **Cost-effective** treatment of internal and external parasites of cattle and pigs.
- ✓ **Low irritant formula.**
- ✓ **Plastic packaging** eliminating the risk of glass breakage.
- ✓ **Convenient** pack size of 500 mL.
- ✓ **Broach period** is 28 days reducing the chance the product will be wasted.
- ✓ **Australian made** which supports local jobs.

CATTLE PARASITES TREATED:

Internal Parasites of Cattle

Internal Parasites – Cattle	Scientific name	Adult worms	Immature worms	Persistent protection*
SMALL BROWN STOMACH WORM	<i>Ostertagia ostertagi</i>	✓	✓ Stage 4 larvae (inc. inhibited larvae)	Up to 21 days
BARBER'S POLE WORM	<i>Haemonchus placei</i>	✓	✓ Stage 4 larvae	Up to 21 days
STOMACH HAIR WORM	<i>Trichostrongylus axei</i>	✓	✓ Stage 4 larvae	Up to 21 days
BLACK SCOUR WORM	<i>Trichostrongylus colubriformis</i>	✓	✓ Stage 4 larvae	
	<i>Trichostrongylus longispicularis</i>	✓		
SMALL INTESTINAL WORM	<i>Cooperia oncophora</i>	✓	✓ Stage 4 larvae	Up to 14 days
	<i>Cooperia</i> spp.	✓	✓ Stage 4 larvae	Up to 21 days
HOOKWORM	<i>Bunostomum phlebotomum</i>	✓		Up to 21 days
NODULE WORM	<i>Oesophagostomum radiatum</i>	✓	✓ Stage 4 larvae	Up to 21 days
(Large) ROUND WORM	<i>Toxocara vitulorum</i>	✓		
LUNGWORM	<i>Dictyoaulus viviparus</i>	✓	✓	Up to 28 days

* Period of persistent protection against reinfection following treatment.

External Parasites of Cattle

External Parasites – Cattle	Scientific name	Activity
CATTLE TICK	<i>Rhipicephalus (Boophilus) microplus</i>	The persistence of Doramate prevents the development of viable cattle ticks for a period of 28 days.
SUCKING LICE	<i>Linognathus vituli</i> <i>Haematopinus eurysternus</i> <i>Solenopotes capillatus</i>	Effective in the treatment of three species of sucking lice in cattle.
MANGE MITE	<i>Chorioptes bovis</i>	Effective in the treatment and control of both <i>Chorioptes bovis</i> and <i>Sarcoptes scabiei</i> var <i>bovis</i> . (Not found in Australia)
BITING LICE	<i>Bovicola bovis</i>	Doramectin aids in the control of biting lice.

AU PRODUCT INFORMATION SHEET



HOW TO USE TROY DORAMATE INJECTABLE ENDECTOCIDE?

Dose rate:

Cattle up to 650 kg		1 mL/ 50 kg body weight
Cattle over 650 kg		1 mL/50 kg body weight
Cattle less than 40 kg		Generally not warranted
Pigs		1 mL/ 33 kg body weight (0.3 mL / 10 kg)
Piglets	5 - 7 kg	0.2 mL
	8 – 10 kg	0.3 mL
	11 – 13 kg	0.4 mL
	14 – 16 kg	0.5 mL

Administration Route:

S.C. in cattle

I.M in pigs

At temperatures below 5 °C
product viscosity increases.
Syringability can be improved
by gently warming the injecting
equipment and product to 15 °C.

Safety: Troy Doramate Injectable Endectocide has a wide safety margin and is safe to administer to cattle and pigs of all ages, including young calves and at all stages of the reproductive cycle in pigs. This product has been specifically formulated to minimise the risk of injection site pain after administration.

Precautions: To avoid tissue residues, take care to estimate the weights and dosages in young animals.

Cattle Tick Control: The persistence of doramectin prevents the development of viable cattle ticks for a period of 28 days following treatment. However, engorged female ticks that drop off up to 3 days after treatment may lay viable eggs and this should be taken into account when planning a strategic tick control program. If presenting cattle for clearance to tick-free areas, consult your local tick advisory authority before use.

Restraint: Do not use in cattle during lactation or less than 60 days before calving where milk or milk products are to be used for human consumption or processing. Do not retreat within 28 days (4 weeks) of the previous treatment.

Withholding Period:

Meat: Cattle: 42 days, Pigs: 35 days.

Milk: Do not use in cattle during lactation or less than 60 days before calving where milk or milk products are to be used for human consumption or processing.

Trade Advice - Export Slaughter Interval (ESI): Cattle and pigs: 42 days.

Storage: Store below 30 °C (room temperature). Store upright.

Please read carton and/or leaflet for full product information.