

AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only



WHAT IS ILIUM TOLFEJEC ANTI-INFLAMMATORY INJECTION?

Ilium Tolfejec is a non-steroidal anti-inflammatory analgesic-antipyretic for use in cattle and pigs.

Physical Description: A clear, colourless/pale yellow solution free of visible particulates.

Schedule: S4

Active Ingredient: 40 mg/mL TOLFENAMIC ACID

WHY CHOOSE ILIUM TOLFEJEC ANTI-INFLAMMATORY INJECTION?

- ✓ **Cost-effective aid** in the treatment of pneumonia and acute mastitis in cattle and metritis-mastitis-agalactia in pigs.
- ✓ **Convenient pack size** of 100 mL.
- ✓ **Broach period is six months** which allows for plenty of time to use the product and reduces the chance that product will be wasted.
- ✓ **Australian made** which supports local jobs.





AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only



HOW TO USE ILIUM TOLFEJEC ANTI-INFLAMMATORY INJECTION?

Dose Rate:

 CATTLE	Pneumonia: 2 mg/kg (1 mL per 20 kg b.w.) by intramuscular injection high in the neck (see leaflet for diagram). Treatment may be repeated once only after 48 hours. Mastitis: 4 mg/kg (1 mL per 10 kg b.w.) as a single intravenous injection.
 PIGS	Metritis-mastitis-agalactia: 2 mg/kg (1 mL per 20 kg b.w.) as a single intramuscular injection.

Route of administration:

I.M. or I.V. dependent on disease.

Restraint:

DO NOT inject cattle other than into muscle tissue high on the side of the neck (see product leaflet for diagram). Injection of product into muscles other than as described is likely to result in residues in meat above the MRL.

Precautions:

Repeat treatments or higher doses could result in residues above the MRL unless the label withholding period is extended. The prescribing veterinarian would need to advise on an extended withholding period. Safe use of ilium Tolfejec Injection during pregnancy has not been established.

Withholding Periods:

Meat: Cattle I.M. administration 10 days, I.V. administration 4 days. Pigs I.M. administration 6 days.

Milk: Cattle I.M. or I.V. administration 12 hours (1 milking). This milk should not be fed to bobby calves.

Trade Advice - Export Slaughter Interval (ESI):

An ESI has not been established for this product.

Storage:

Store below 30 °C (room temperature). Protect from light.

Please refer to carton and/or enclosed leaflet for full product information.

