


AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

**ilium
Tula**
INJECTION



WHAT IS ILIUM TULA INJECTION?

ilium Tula Injection is a tulathromycin antibiotic for the treatment of respiratory infections in cattle and pigs.

Active Ingredient: 100 mg/mL tulathromycin

Poison Schedule: S4

Physical Description: A clear, colourless to slightly yellow solution.

WHY CHOOSE ILIUM TULA INJECTION?

✓ **Cost-effective** treatment of:

- Bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* sensitive to tulathromycin.
- Swine respiratory disease (SRD) associated with *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, and *Haemophilus parasuis* sensitive to tulathromycin.

✓ **Handy 100 mL pack size** to maximise usability.

✓ **28 day broach period** which allows time to use the product and reduces the chance of wastage.

✓ **Premium syringability** due to the formulation's low viscosity.

✓ **Australian made.**



www.troylab.com.au



(+61) 2 8808 3600




sales@troylab.com.au

TROY

AU PRODUCT INFORMATION SHEET



Information Provided for Veterinary Professionals Only

**ilium
Tula**
INJECTION



HOW TO USE ILIUM TULA INJECTION?

Dose Rates:

ANIMAL	DOSE RATE	ADMINISTRATION ROUTE
 CATTLE	1 mL/40 kg body weight with no more than 10 mL injected at one site.	Cattle: single subcutaneous injection high on the neck.
 PIGS	1 mL/40 kg body weight with no more than 2.5 mL injected at one site.	Pigs: single intramuscular injection in the neck.

Restraints:

Use only in respiratory infections of cattle and swine.

DO NOT USE in dairy cows (except replacement dairy heifers) that are producing milk or may in the future produce milk for human consumption or processing.

Do not use in bobby calves.

Do not treat dairy heifers following the first mating.

Side Effects:

Cattle: S.C. injection frequently causes transient pain reactions and local swellings at the injection site that can persist for up to 30 days.

Pigs: Pain reactions have not been observed in pigs after I.M. injection, but local swellings at the injection site can persist up to 30 days. Mild salivation that resolves in less than 4 hours have also been reported.

Re-Treatment Intervals:

Cattle: Do not re-treat cattle for 12 weeks after last treatment. Dairy heifers: Do not re-treat dairy heifers.

Pigs: Do not re-treat pigs for 8 weeks after last treatment.

Withholding Periods:

- MEAT: Cattle: 35 days. Pigs: 14 days.
- MILK: **DO NOT USE** in dairy cows (except replacement dairy heifers) that are producing milk or may in the future produce milk for human consumption or processing.

Trade Advice - Export Slaughter Interval (ESI):

Cattle: 35 days. **Pigs:** 26 days.

Storage: Store below 30 °C (room temperature).

Please refer to carton and/or enclosed leaflet for full product information.

